

Improved pharmacy experience for patients

Smith+Nephew

Collagenase
SANTYL[®]
Ointment 250 units/gram

- Local:** Over 25,000 pharmacy locations
Convenient: Rx pick-up*
Accurate: Proper dosing for each patient
- Option 1 - Provide the following:**
- # of days therapy
 - Instructions for use
 - The following must be entered in the “Notes to Pharmacy” section of the e-prescription.
 - Wound dimensions
 - Dispense quantity sufficient per manufacturer’s dosing calculator”

Collagenase SANTYL[®] Ointment (250 units/g)
Apply to wound as directed once daily for days
Wound Dim 1: cm X cm
Wound Dim 2: cm X cm
Wound Dim 3: cm X cm
Wound Dim 4: cm X cm

Dispense quantity sufficient per manufacturer’s dosing calculator

REFILLS

<https://santyl.com/hcp/dosing>



- Option 2**
When necessary information is missing, pharmacy team calls within 24 business hours to obtain details for correct dosing.

Participating pharmacies



The Kroger Family of Pharmacies



Important Safety Information:

Indications: Collagenase SANTYL Ointment (“SANTYL”) is indicated for debriding chronic dermal ulcers and severely burned areas.

Contraindications: SANTYL is contraindicated in patients who have shown local or systemic hypersensitivity to collagenase.

Warnings and Precautions: The optimal pH range of collagenase is 6 to 8. Higher or lower pH conditions will decrease the enzyme’s activity and appropriate precautions should be taken. The enzymatic activity is also adversely affected by certain detergents, and heavy metal ions such as mercury and silver which are used in some antiseptics. As such, the wound should be properly cleansed prior to application of SANTYL. Debilitated patients should be closely monitored for systemic bacterial infections because of the theoretical possibility that debriding enzymes may increase the risk of bacteremia. A slight transient erythema has been noted occasionally in the surrounding tissue, particularly when SANTYL was not confined to the wound. SANTYL is not indicated for wound closure. Discontinue use of SANTYL after granulation tissue is well-established.

Adverse Reactions: No allergic sensitivity or toxic reactions have been noted in clinical use when used as directed. The risk information provided herein is not comprehensive. To see the complete Prescribing Information, please see the FDA-approved product labeling, here: <https://www.santyl.com/pdf/SANTYL-PI.pdf>. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch or call 1-800-FDA-1088.

Advanced Wound Management

Smith+Nephew, Inc.
Fort Worth, TX 76109
USA

www.smith-nephew.com
www.santyl.com

Customer Care Center

T 800-876-1261
F 727-392-6914

[®]Trademark of Smith+Nephew
[™]All trademarks acknowledged
© 2023 Smith+Nephew, Inc.
SAPE27-38438-0123

